

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human S100A13 in ELISAs and Western blots. In direct ELISAs, approximately 50% cross-reactivity with recombinant mouse S100A13 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 854114
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human S100A13 Ala2-Lys98 Accession # Q99584
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

S100A13 is an 11 kDa member of the S100 (soluble in 100% saturated ammonium sulfate) family of vertebrate EF-hand Ca⁺⁺-binding proteins (1-3). It is widely expressed as a homodimer with two 98 amino acid (aa) long subunits (2, 3). Human S100A13 shares 83%, 90%, 91%, 87%, 78% and 47% aa identity with mouse, rat, cow, dog, opossum and chicken S100A13, respectively. Like other S100 proteins, S100A13 is small and generally acidic, but contains a basic residue-rich sequence at the C terminus, and two EF hand motifs that bind with Ca⁺⁺ differing affinities (2-4). Some S100 proteins, including S100A13, are able to bind the cell surface receptor for advanced glycation end-products (RAGE) (5). Despite lacking a signal sequence, S100A13 plays an important role in Cu⁺⁺-dependent export of FGF-1 (FGF acidic) and IL-1α from the cell in response to stresses such as heat shock, anoxia and starvation (6-8). Binding of copper is necessary for formation of a multi-protein complex between S100A13, FGF-1 and p40 synaptotagmin-1 (synt-1) (9, 10). Cu⁺⁺ ions supplied by S100A13 are thought to oxidize and downregulate the activity of FGF-1 prior to export (10). Calcium influx may also play a similar role in FGF-1 release from neuronal cells (11). S100A13 is composed of four amphiphilic helices that may interact with acidic phospholipid headgroups. With FGF-1 and synt-1, S100A13 likely perturbs the membrane, which allows the S100A13 protein complex to exit the cell (4, 12). S100A13 has been proposed as a marker for angiogenesis in tumors and endometrium, due to its role in stress-induced export of FGF-1 (13, 14). Based on in house studies, S100A13 has also been found to promote neurite outgrowth from rat cortical embryonic neurons (15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.