

DESCRIPTION

Species Reactivity	Human
Specificity	Detects endogenous human PON2 in Western blots. In Western blots, this antibody does not cross-react with recombinant human (rh) PON1 or rhPON3.
Source	Monoclonal Rat IgG _{2A} Clone # 453709
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human PON2 Ala30-Leu354 Accession # NP_000296
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The paraoxonase (PON) gene family of antioxidant enzymes includes three known members located adjacent to each other on chromosome 7. Paraoxonase/arylesterase 2 (PON2) is a 354 amino acid, 39 kDa protein that is widely expressed in a variety of tissues and may act as a cellular antioxidant, protecting cells from oxidative stress. PON2 has arylesterase and arylalkylphosphatase activity (EC 3.1.1.2 and EC 3.1.8.1) and can hydrolyze a number of organophosphate substrates and aromatic carboxylic acid esters. PON2 is membrane-bound and has several potential glycosylation sites. Sequence polymorphisms in this gene may be associated with coronary heart disease and a number of phenotypes related to diabetes. PON2 is not associated with HDL but can prevent LDL lipid peroxidation and reverse the oxidation of mildly oxidized LDL. Alternatively spliced transcript variants encoding different isoforms have been described.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.