

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse TRAM/TICAM2 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) TRAM/TICAM2 or rhTRIF/TICAM1 is observed.
Source	Monoclonal Rat IgG _{2B} Clone # 757706
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse TRAM/TICAM2 Met1-Ala232 Accession # Q8BJQ4
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The innate and adaptive immune responses depend on systems that link cell surface surveillance receptor signals to cytoplasmic proteins such as kinases, adaptors, and transcription factors. Toll-like receptors (TLR) recognize different pathogen associated molecular patterns (PAMPs), and initiate a signaling cascades mediated by a Toll/interleukin-1 receptor (TIR) domain-containing adaptor proteins such as MyD88, TIRAP/MAL, and TRIF. Mouse TRIF-related adaptor molecule (TRAM), is a 232 amino acid, 26 kDa (predicted), ubiquitously expressed member of the TIR domain-containing adaptor family. TRAM, also known as TIR domain-containing adapter protein 2 (TICAM2) and TIR domain-containing protein (TIRP), contains a central TIR domain that is most similar to that of TRIF. TRAM plays an essential role in the MyD88-independent signaling of TLR4 by binding members of the IRAK family, ultimately leading to the activation of NFκB. Mouse TRAM shares 75% and 77% identity to human and rat TRAM, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.