

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SIRPy/CD172g in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 20% cross-reactivity with recombinant human (rh) SIRPα and rhSIRPβ 1 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 595337
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human SIRPy/CD172g Val64-Ser364 Accession # Q9P1W8
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Signal-regulatory protein gamma/beta 2 (SIRP gamma/beta 2) is a 45-47 kDa type 1 transmembrane glycoprotein and member of the immunoglobulin superfamily and SIRP family of proteins. Human SIRP gamma/beta 2 is synthesized as a 387 amino acid (aa) precursor that contains a 28 aa signal sequence, a 332 aa extracellular domain (ECD), a 23 aa transmembrane segment, and a 4 aa cytoplasmic tail. The ECD contains two Ig-like C1-type domains, one Ig-like V-type domain, and four potential sites for N-linked glycosylation. There are four named isoforms for human SIRP gamma/beta 2. Human SIRP gamma/beta 2 has no orthologs, but it is 75% aa identical to SIRPβ. It is expressed in the liver and at very low levels in brain, heart, lung, pancreas, kidney, placenta, and skeletal muscle.

SIRP gamma/beta 2 is also expressed on CD4⁺ T-cells, CD8⁺ T-cells, and activated NK cells.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.