

**DESCRIPTION**


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse IL-17 R in direct ELISAs and Western blots. In Western blots, approximately 5% cross-reactivity with recombinant human IL-17 R is observed and less than 1% cross-reactivity with recombinant mouse (rm) IL-17 RC and rml-17 RD is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant mouse IL-17 R Extracellular domain
<b>Conjugate</b>	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	10 $\mu$ L/10 <sup>6</sup> cells	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

**BACKGROUND**

IL-17 R, also known as IL-17 RA, is a 120 kDa type I transmembrane glycoprotein protein that plays a central role in inflammatory responses (1-3). Mature mouse IL-17 R consists of a 291 amino acid (aa) extracellular domain, a 21 aa transmembrane segment, and a 521 aa cytoplasmic domain (4). The cytoplasmic domain contains a region homologous to the TIR domain of the TLR/IL-1 R family (5). Mouse IL-17 R shares 84% and 72% aa sequence identity with rat and human IL-17 R, respectively. Within the extracellular domain, it shares 18-25% sequence identity with mouse IL-17 RB, C, D, and E. While the expression of IL-17 is restricted to activated T cells, IL-17 R exhibits a broad tissue distribution (4). Even in the absence of ligand, IL-17 R exists on the cell surface as a multimer (6). IL-17 R can bind IL-17 but must associate with IL-17 RC to transduce signals (7, 8). Interestingly, human IL-17 R does not appear to form productive complexes with mouse IL-17 RC (8). The IL-17 R can also signal in response to IL-17F (9). IL-17 R ligation promotes T cell activation and the production of IL-6, G-CSF, SCF, and multiple pro-inflammatory chemokines (4, 7, 9, 10). IL-17A and IL-17F synergize with TNF- $\alpha$  in the induction of CXCL1, G-CSF, and IL-6 (9, 11). This effect requires the presence of both TNF RI and TNF RII (9). IL-17 interactions with IL-17 R also inhibit the TNF- $\alpha$  induced upregulation of fibroblast CCL5 and VCAM-1 (11). CCL5 and VCAM-1 induced effects are differentially sensitive to blockade with IL-17 R specific antibodies, suggesting that IL-17 R triggers divergent intracellular signals (11). *In vivo*, IL-17 R activity is important for increased generation of neutrophils and their recruitment to sites of inflammation (10, 12, 13). IL-17 R is required for host defense against microbial infection and for the progression of arthritis from inflammation to destructive joint erosion (10, 13).

**References:**

1. Iwakura, Y. and H. Ishigame (2006) *J. Clin. Invest.* **116**:1218.
2. Moseley, T.A. *et al.* (2003) *Cytokine Growth Factor Rev.* **14**:155.
3. Kawaguchi, M. *et al.* (2004) *J. Allergy Clin. Immunol.* **114**:1265.
4. Yao, Z. *et al.* (1995) *Immunity* **3**:811.
5. Novatchkova, M. *et al.* (2003) *Trends Biochem. Sci.* **28**:226.
6. Kramer, J.M. *et al.* (2006) *J. Immunol.* **176**:711.
7. Hymowitz, S.G. *et al.* (2001) *EMBO J.* **20**:5332.
8. Toy, D. *et al.* (2006) *J. Immunol.* **177**:36.
9. McAllister, F. *et al.* (2005) *J. Immunol.* **175**:404.
10. Ye, P. *et al.* (2001) *J. Exp. Med.* **194**:519.
11. Schnyder, B. *et al.* (2005) *Cytokine* **31**:191.
12. Tan, W. *et al.* (2006) *J. Immunol.* **176**:6186.
13. Lubberts, E. *et al.* (2005) *J. Immunol.* **175**:3360.