

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse LIF in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant mouse (rm) IL-6, rmIL-11, recombinant human (rh) LIF, rmCT-1, rhCLC or recombinant rat CNTF is observed.
Source	Monoclonal Rat IgG _{2B} Clone # 139124
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse LIF Pro25-Phe203
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Leukemia inhibitory factor (LIF) was initially identified as a factor that inhibited the proliferation and induced the differentiation to macrophages of the murine myeloid leukemic cell line M1. Subsequent to its purification and molecular cloning, LIF was recognized to be a pleiotropic factor with multiple effects on both hematopoietic and non-hematopoietic cells. LIF has overlapping biological functions with OSM, IL-6, IL-11 and CNTF. All these cytokines utilize gp130 as a component in their signal transducing receptor complexes. Mouse LIF cDNA encodes a 203 amino acid residue polypeptide with a 23 amino acid signal peptide that is cleaved to yield a 180 amino acid mature mouse LIF. Native human and mouse LIF are highly glycosylated monomeric proteins. Both human and murine LIF protein sequences have multiple potential N- and O-linked glycosylation sites and six conserved cysteine residues that are involved in three intramolecular disulfide bridges.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.