

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CCL3/MIP-1 α in direct ELISAs.
Source	Monoclonal Rat IgG ₁ Clone # 756605
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse CCL3/MIP-1 α Ala24-Ala92 Accession # P10855
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CCL3, also known as Macrophage Inflammatory Protein 1 alpha (MIP-1 α) and LD78, is a member of the β or CC subfamily of chemokines and is closely related to CCL4/MIP-1 β . Chemokines comprise a large family of small secreted proteins that are involved in immune and inflammatory responses. CCL3 expression can be induced in a variety of hematopoietic cells, fibroblasts, smooth muscle cells, and epithelial cells (1). Mature mouse CCL3 shares 73%, 91%, and 82% amino acid sequence identity with human, rat, and cotton rat CCL3, respectively (2). CCL3 is an approximately 8 kDa chemokine that forms complexes with sulfated proteoglycans (3, 4). In a reversible process, CCL3 associates into noncovalently-linked dimers which then form tetramers and high molecular weight polymers (5, 6). These complexes of CCL3 are protected from proteolytic digestion by Insulin Degrading Enzyme (IDE) which can cleave the monomeric chemokine (6). CCL3 exerts its biological functions through interactions with CCR1, CCR3, and CCR5 (1). It is cleared from the extracellular space by internalization via the decoy chemokine receptor D6 (7). CCL3 promotes the chemoattraction, adhesion to activated vascular endothelium, and cellular activation of many hematopoietic cell types including activated T cells, NK cells, neutrophils, monocytes, immature dendritic cells, and eosinophils (1, 8-10). CCL3 is also known as Stem Cell Inhibitor (SCI) and can inhibit the proliferation of hematopoietic progenitor cells (3). CCL3 bioactivity contributes to tumor metastasis and the inflammatory components of viral infection, rheumatoid arthritis, and hepatitis (11-14), although it also can suppress the replication of HIV (15). CCL3 additionally promotes hyperalgesia by sensitizing sensory neurons to TRPV1-mediated noxious stimulation (16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.