

#### DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human CD81 in Elisas.
<b>Source</b>	Monoclonal Rabbit IgG Clone # 2664C
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	CHO-derived human CD81 protein Phe113-Ala201 Accession # P60033
<b>Conjugate</b>	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

**Western Blot** Optimal dilution of this antibody should be experimentally determined.

#### PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

#### BACKGROUND

CD81, also known as TAPA-1 and Tetraspanin-28, is an approximately 25 kDa palmitoylated component of plasma membrane lipid rafts (1). It contains four transmembrane segments, two extracellular loops of 30 and 90 amino acids (aa), and three short cytoplasmic regions (2, 3). Within the large extracellular loop, human CD81 shares 90% and 84% aa sequence identity with mouse and rat CD81, respectively. CD81 associates with a wide range of membrane proteins including CD151, TfR2, LDL R, PCSK9, Glypican 3, IFITM1, IGSF8/CD316, FPRP, and complexes of CD19-CD21 (4-11). It is required for the development of CD4+CD8+ DP thymocytes (12) and hepatocyte infection by Plasmodium sporozoites (13). It also supports B cell receptor signaling (11), Hepcidin expression (5), monocyte and B cell tethering to the vascular endothelium (14), and the immunosuppressive function of Treg and MDSC (15). CD81 additionally functions as a receptor for the E2 glycoprotein of hepatitis C virus (16). The CD81-E2 interaction inhibits NK cell cytolytic activity, provides a co-stimulatory signal to T cells, and inhibits the maturation of plasmacytoid dendritic cells (17-19).

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.