

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CRACC/SLAMF7 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human CRACC, recombinant mouse (rm) CD229, rmCD84, or rmNTB-A is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 520911
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant mouse CRACC/SLAMF7 Ala22-Gly224 Accession # Q8BHK6
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD2-like receptor activating cytotoxic cells (CRACC), also known as CS1, novel Ly9, SLAMF7, and CD319, is a 66 kDa type I transmembrane glycoprotein in the SLAM subgroup of the CD2 family (1). Mature mouse CRACC consists of a 202 amino acid (aa) extracellular domain (ECD) with one Ig-like V-set domain and one Ig-like C2-set domain, a 21 aa transmembrane segment, and an 88 aa cytoplasmic domain with two immunoreceptor tyrosine-based switch motifs ITSMs (2, 3). Within the ECD, mouse CRACC shares 53% aa sequence identity with human CRACC. It shares 19%-35% aa sequence identity with comparable regions of other mouse SLAM proteins including 2B4, BLAME, CD2F-10, CD84, CD229, NTB-A, and SLAM/CD150. Additional isoforms of mouse CRACC are distinguished by deletions and/or substitutions in their cytoplasmic domains. CRACC is expressed on the surface of NK cells, CD8⁺ T cells, activated B cells, and mature dendritic cells (4, 5). It interacts homophilically to induce NK, CTL, and B cell activation (4-7). In human NK cells, activated CRACC transmits signals following association with the adaptor protein EAT-2 (8).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.