

#### DESCRIPTION

|                           |                                                                                                                                                                                                                                                                             |
|---------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b> | Mouse                                                                                                                                                                                                                                                                       |
| <b>Specificity</b>        | Detects mouse CCL9/10/MIP-1 $\gamma$ in ELISAs and Western blots. In sandwich ELISAs, less than 0.02% cross-reactivity with recombinant human CCL3, 4, 15, 19, 20, recombinant mouse CCL3, 4, 19, or recombinant rat CCL20 is observed.                                     |
| <b>Source</b>             | Monoclonal Rat IgG <sub>1</sub> Clone # 62105                                                                                                                                                                                                                               |
| <b>Purification</b>       | Protein A or G purified from hybridoma culture supernatant                                                                                                                                                                                                                  |
| <b>Immunogen</b>          | <i>E. coli</i> -derived recombinant mouse CCL9/10/MIP-1 $\gamma$<br>Gln22-Gln122<br>Accession # P51670                                                                                                                                                                      |
| <b>Conjugate</b>          | Alexa Fluor 750<br>Excitation Wavelength: 749 nm<br>Emission Wavelength: 775 nm                                                                                                                                                                                             |
| <b>Formulation</b>        | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

|                                                |                                                                        |
|------------------------------------------------|------------------------------------------------------------------------|
| <b>ELISA Capture (Matched Antibody Pair)</b>   | Optimal dilution of this antibody should be experimentally determined. |
| <b>ELISA Detection (Matched Antibody Pair)</b> | Optimal dilution of this antibody should be experimentally determined. |
| <b>Neutralization</b>                          | Optimal dilution of this antibody should be experimentally determined. |
| <b>Western Blot</b>                            | Optimal dilution of this antibody should be experimentally determined. |

#### PREPARATION AND STORAGE

|                                |                                                                                                                   |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

#### BACKGROUND

Mouse CCL9/10 (also named MIP-1 $\gamma$  and MRP-2) is an 11 kDa, secreted, monomeric polypeptide that belongs to the  $\beta$  (or CC) intercrine family of chemokines (1-3). Based on its activity and amino acid (aa) sequence, it is further classified as a member of the NC6 or six cysteine-containing CC subfamily of chemokines (2, 4, 5). This subfamily contains four N-terminally extended chemokines, two human (CCL15 and CCL23) and two mouse (CCL9 and CCL10). Within this subfamily, there are no human-to-rodent interspecies orthologs. Mouse CCL9/10 is synthesized as a 122 aa precursor that contains a 21 aa signal sequence and a 101 aa mature region with six cysteines. As noted, the mature region has an expanded N-terminus relative to other CC family members, and it forms a third intrachain disulfide bond with its two extra cysteines (3-7). Mouse CCL9/10 is 75% aa identical to rat CCL9/10 (8). Chemokines are known to undergo proteolytic processing to generate multiple isoforms. NC6 chemokines are usually only marginally active at full-length, but are converted to highly active forms upon N-terminal truncation. Mature CCL9, in the presence of inflammatory fluids, is naturally truncated by 28, 29 or 30 aa at the N-terminus, generating a highly active, 8 kDa, 71-73 aa CCR1 ligand. In contrast, other CCR1 ligands, CCL3/MIP-1 $\alpha$  and CCL5/RANTES, lose their potency when proteolytically processed. CCL9/10 is constitutively secreted, and circulates as a full-length molecule. Any onset of inflammation with subsequent enzyme release may act on local NC6 chemokines, generating early, potent leukocyte chemoattractants (5, 7).

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.