

DESCRIPTION

Species Reactivity	Human
Specificity	Detects endogenous human GRAP2 in Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 475804
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human GRAP2 Met1-Arg330 Accession # O75791
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

GRAP2 (GRB2 related adaptor protein 2), a divergent member of the GRB2/Sem5/Drk family, is an adaptor protein involved in leukocyte-specific protein-tyrosine kinase signaling. GRAP2 shares similar structural organization as GRB2 and GRAP with a central SH2 domain flanked by 2 SH3 domains. However, unlike GRB2 and GRAP, the SH2 domain and the C-terminal SH3 domain of GRAP2 contains a unique 120-amino acid glutamine/proline rich sequence. In T lymphocytes, GRAP2 interacts with LAT and SLP-76 to regulate NF-AT (nuclear factor of activated T cells) and form a signaling complex with HPK1 to mediate the JNK signaling pathway. GRAP2 also interacts with M-CSF R and the activated T cell co-stimulatory receptor CD28.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.