

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human NAALADase-like 2/NAALADL2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 817225
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human NAALADase-like 2/NAALADL2 Ser152-Asn795 Accession # Q58DX5
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the [Technical Information](#) section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

NAALADL2 (N-acetylated α -linked acidic dipeptidase like 2) is a member of the peptidase M28 family of enzymes. It is presumably a type II transmembrane (TM) protein that may have O-glycosyl hydrolase activity. Human NAALADL2 is 795 amino acids (aa) in length. It is believed to contain a cytoplasmic and TM segment at the N-terminus, followed by a peptidase domain (aa 444-596) and a TfR-like dimerization region (aa 688-777). There are multiple splice forms. One shows a 4 aa substitution for aa 292-795, a second shows an alternate start site at Met18 with a 24 aa substitution for aa 314-795, and a third contains an alternate start site at Met283, accompanied by a deletion of aa 363-411 and 633-795. Over aa 152-795, human NAADADL2 shares 87% and 82% aa sequence identity with mouse and canine NAADADL2, respectively.