

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CCL2 in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 123616R
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant mouse CCL2 Gln24-Arg96 Accession # P10148
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	Mouse splenocytes treated with LPS and monensin

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Mouse CCL2 is a member of the β (C-C) subfamily of chemokines. The mouse CCL2 gene was initially identified as a platelet-derived growth factor-inducible gene in mouse fibroblasts. Mouse CCL2 cDNA encodes a 148 amino acid (aa) residue with a putative 23 aa signal peptide that is cleaved to generate the mature protein. Mouse CCL2 shares 82% amino acid sequence identity with rat CCL2. Mouse CCL2 also shares 55% amino acid sequence identity with human MCP-1. Compared to human MCP-1, mouse CCL2 has a 49 aa residue extension at the carboxy-terminus. When a DNA sequence encoding the 125 aa residue of the mature CCL2 protein was expressed in *E. coli* at R&D Systems, the purified protein had the predicted N-terminus but a mass of 8525 Da. The truncation of most of the C-terminal extension could be due either to purification artifact or to post-translational modification. The truncated recombinant CCL2 has a potency similar to that of human MCP-1 in the monocyte chemotaxis assay. Mouse CCL2 has full activity on human cells while human MCP-1 has limited activity on mouse cells.

References:

1. Rollins, B.J. *et al.* (1988) *Proc. Natl. Acad. Sci. USA* **85**:3738.
2. Gu, L. *et al.* (1999) *Chem. Immunol.* **72**:7.
3. Luini, W. *et al.* (1994) *Cytokine* **6**:28.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.