

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human HLA-DR.
Source	Monoclonal Mouse IgG ₁ Clone # L203
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	RPMI 8866 human lymphoblastoid cells Accession # P01903
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 µL/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

HLA-DR is a transmembrane human major histocompatibility complex 2 (MHC II) family member and consists of a 34 kDa (alpha) subunit and one of several 28 kDa (beta) subunits. HLA-DR is expressed primarily by B cells and dendritic cells (DC), in which it binds peptides derived from internalized and processed antigenic proteins. It presents these peptides on the cell surface for recognition by the T cell receptor on CD4⁺ T cells. This interaction is central to antigen specificity in the adaptive immune response. HLA-DR alleles, polymorphisms, and aberrant expression are linked to a variety of diseases including autoimmunity and cancer.