

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human NGL-1/LRRC4C in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) LRRC4 or rhNGL-3/LRRC4B is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 755720
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human NGL-1/LRRC4C Gln45-Lys527 (predicted) Accession # Q9HCJ2
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human NGL-1 (Netrin-G1 ligand; gene name LRRC4C) is a 67 kDa (predicted for mature protein), type I transmembrane cell adhesion molecule that is a member of the NGL family of proteins (1, 2). It is synthesized from a precursor that is 640 amino acids (aa) in length that contains a 44 aa signal sequence, a 483 aa extracellular region, a 21 aa transmembrane region, and a short cytoplasmic tail of 92 aa. The extracellular region of NGL-1 consists of nine leucine-rich repeats (LRRs) that are flanked by LRR N-terminal and LRR C-terminal domains, and followed by an Ig-like C2-type domain (1, 2). The cytoplasmic region contains a C-terminal Glu-Thr-Gln-Ile sequence that corresponds to a potential PDZ (postsynaptic density-95/discs large/zona occludens-1) domain-binding motif (1, 2). Human NGL-1 shares 99.7% aa sequence identity with mouse NGL-1. Mouse NGL-1 is highly expressed in the developing cerebral cortex and the striatum at embryonic day 14 (1). Postnatally, NGL-1 is expressed exclusively in the brain, with the highest expression found in the cerebral cortex as a whole, and in individual neocortical areas such as the frontal, parietal and occipital lobes (1). Moderate expression of NGL-1 occurs in the putamen, amygdala, hippocampus and medulla oblongata (1). Weak expression is found in the caudate nucleus and thalamus (1). Functionally, membrane-bound cell-surface NGL-1 binds to netrin-G1 specifically through its LRR region, and in the developing brain, may promote neurite outgrowth of thalamocortical axons (1-4).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.