

## Human Siglec-14 Alexa Fluor® 405-conjugated Antibody

Recombinant Monoclonal Rabbit IgG Clone # 2457E Catalog Number: FAB4905V 100 µg

DESCRIPTION		
Species Reactivity	Human	
Specificity	Detects human Siglec-14 in direct ELISAs.	
Source	Recombinant Monoclonal Rabbit IgG Clone # 2457E	
Purification	Protein A or G purified from cell culture supernatant	
Immunogen	Mouse myeloma cell line, NS0-derived human Siglec-14 Lys18-Leu358 Accession # Q08ET2	
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm	
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS			
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.			
	Recommended	Sample	
	Concentration		
Flow Cytometry	0.25-1 μg/10 <sup>6</sup> cells	HEK293 Human Cell Line Transfected with Human Siglec-14	

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze.	
	12 months from date of receipt, 2 to 8 °C as supplied.	

## BACKGROUND

Siglecs are sialic acid specific I-type lectins that belong to the immunoglobulin superfamily. Structurally, they are transmembrane proteins with an N-terminal Ig-like V-set domain followed by a varying number of Ig-like C2-set domains (1, 2). Orthologs of Siglec-14 have not been described in mouse or rat. Mature human Siglec-14 consists of a 342 amino acid (aa) extracellular domain (ECD) with one Ig-like V-set domain and two Ig-like C2-set domains, a 23 aa transmembrane segment, and a 15 aa cytoplasmic tail (3). Within the first two Ig-like domains, Siglec-14 shares 99.5% aa sequence identity with Siglec-5, and 51% - 56% with Siglec-3, -6, -7, -8, and -9. Siglec-5 and -14 exhibit similar ligand preference among sialylated glycans, although Siglec-14 binds with higher avidity (3). Siglec-14 does not have the cytoplasmic ITIM sequence that mediates inhibitory signaling of most other Siglecs (2, 3). However, its transmembrane segment contains a charged arginine residue that enables association with the adaptor protein DAP12 (3, 4). Siglec-15 is the only other human Siglec identified to date that associates with DAP12 (5). Siglec-14 mRNA is primarily expressed in bone marrow, spleen, and fetal liver (3). The protein may be expressed on neutrophils, monocytes, and macrophages, as reports describing Siglec-5 on these cells employed some antibodies that also detect Siglec-14 (3, 6, 7). Siglec-5 and -14 likely function as paired receptors with similar ligand specificity and cellular expression but potentially opposing effects on cellular activation (3).

## References:

- 1. Varki, A. and T. Angata (2006) Glycobiology 16:1R.
- 2. Crocker, P.R. et al. (2007) Nat. Rev. Immunol. 7:255.
- 3. Angata, T. et al. (2006) FASEB J. 20:1964.
- 4. Turnbull, I.R. and M. Colonna (2007) Nat. Rev. Immunol. 7:155.
- 5. Angata, T. et al. (2007) Glycobiology 17:838.
- 6. Connolly, N.P. et al. (2002) Br. J. Haematol. 119:221.
- 7. Erickson-Miller, C.L. et al. (2003) Exp. Hematol. 31:382.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 8/1/2019 Page 1 of 1

