

Human Ninjurin-1 Alexa Fluor® 750-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 758926

Catalog Number: FAB51051S

100 µg

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Ninjurin-1 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human Ninjurin-2 is observed.
Source	Monoclonal Mouse IgG ₁ Clone # 758926
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Ninjurin-1 Asp2-Val81 Accession # Q92982
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	HepG2 human hepatocellular carcinoma cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Ninjurin-1 (nerve injury-induced protein 1) is a 20-22 kDa member of the Ninjurin family of transmembrane (TM) proteins. It is expressed by Schwann cells, neurons and hepatocytes and participates in intercellular homophilic binding during nerve regeneration. Human Ninjurin-1 is 152 amino acids in length. It has an unusual membrane orientation. There is an 80 amino acid (aa) N-terminal extracellular domain (ECD) (aa 1-80), followed by a TM segment, a cytoplasmic region, a second TM segment and a C-terminal ECD (aa 142-152). Homophilic binding is divalent-cation dependent and occurs between Pro26 and Asn37. Over aa 1-80, human Ninjurin-1 shares 84% aa sequence identity with mouse Ninjurin-1. Human Ninjurin-1 shares only 50% aa sequence identity with human Ninjurin-2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.