

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human DGK-ε in direct ELISAs and human, mouse and rat DGK-ε in Western blots. In Western blots, approximately 10-50% cross-reactivity with recombinant human DGK zeta, eta, iota, alpha, theta, gamma, kappa, delta, and beta is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 670914
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human DGK-ε Asn314-Arg435 Accession # P52429
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DGK-ε (Diacylglycerol kinase epsilon) is a 65 kDa member of the eukaryotic diacylglycerol kinase family of enzymes. It is a type III DGK that possesses only a C1/Cys-rich domain and a catalytic region, and is found in neurons and testis. DGK-ε specifically phosphorylates arachidonate-containing DAG, and may downregulate DAG signaling that results from inositol cycling. Human DGK-ε is 567 amino acids (aa) in length. It contains one predicted transmembrane domain (aa 22-42), two C1 DAG-binding regions (aa 59-108 and 124-177) and one catalytic domain (aa 219-350). DGK-ε is predicted to form intramembrane oligomers. Over aa 314-435, human DGK-ε shares 99% aa identity with mouse DGK-ε.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.