

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat CXCL1/GRO α /KC/CINC-1 in direct ELISAs.
Source	Monoclonal Rabbit IgG Clone # 2176C
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant rat CXCL1/GRO α /KC/CINC-1 Ala25-Lys96 Accession # P14095
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cytokine-induced Neutrophil Chemoattractant 1 (CINC-1) was originally purified from media conditioned by IL-1 β stimulated rat kidney epithelioid cells (NRK-52E). On the basis of its protein and DNA sequences, CINC-1 is a member of the alpha (CXC) subfamily of chemokines, designated CXCL1. Three additional rat CXC chemokines (CINC-2 α , CINC-2 β , CINC-3/MIP-2), sharing approximately 63-67% amino acid sequence identity with CINC-1, have been identified. The protein sequence of rat CINC-1 is also 68%, 71%, and 69% identical to that of human GRO- α , GRO- β , and GRO- γ , respectively. Based on their sequence homology, it has been suggested that CINC-1 is the rat counterpart of human GROs. CINC-1 is also the counterpart of mouse KC. Rat CINC-1 cDNA encodes a 96 amino acid residue precursor protein from which the amino-terminal 24 amino acid residues are cleaved to generate the mature CINC-1. Similar to other alpha chemokines, rat CINC-1s are potent neutrophil attractants and activators and have been shown to play an important role in the infiltration of neutrophils into inflammatory sites in rats.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.