

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human VSIG2 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 541522
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human VSIG2 Val24-Ala243 Accession # Q961Q7-1
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	HEK293 Human Cell Line Transfected with Human VSIG2 and eGFP

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

V-set and immunoglobulin domain-containing protein 2 (VSIG2), also known as cortical thymocyte-like protein (CT-like protein), is a member of the immunoglobulin superfamily (IgSF) that was originally described in *Xenopus* and was referred to as cortical thymocyte marker in *Xenopus*, or CTX. The clone isolated from the human genomic library was called CTH (1). Mature human VSIG2 is a single-pass type I transmembrane protein with a 220 amino acid (aa) extracellular domain (ECD) that contains one C2-type and one V-type immunoglobulin-like domains, and a 63 aa cytoplasmic region linked to the ECD by a short 21 aa transmembrane domain. Human VSIG2 ECD shares 85% and 86% aa sequence identity with mouse and rat VSIG2, respectively. VSIG2 is structurally related to the B7 family of immune regulatory proteins. Our studies at R&D Systems show that VSIG2 inhibits T cell activation, including IL17 and interferon gamma production.

References:

1. Chrétien, I. *et al.* (1998) Eur. J. Immunol. **28**:4094.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.