

**DESCRIPTION**

<b>Species Reactivity</b>	Rat
<b>Specificity</b>	Detects rat IL-18/IL-1F4 in direct ELISAs and Western blots. In direct ELISAs, this antibody shows 100% cross-reactivity with recombinant mouse IL-18 and no cross-reactivity with recombinant human IL-18, recombinant porcine IL-18, recombinant rat (rr) IL-18.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 69604
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant rat IL-18/IL-1F4 His37-Ser194 Accession # P97636.1
<b>Conjugate</b>	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
<b>Formulation</b>	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide

\*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the Technical Information section on our website.

<b>Western Blot</b>	Optimal dilution of this antibody should be experimentally determined.
---------------------	--

**PREPARATION AND STORAGE**

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

**BACKGROUND**

Interleukin-18 (IL-18), also known as IL-1F4 and IFN- $\gamma$  inducing factor (IGIF), is a member of the IL-1 family of cytokines and is a key molecule in the innate immune response (1). Rat IL-18 is synthesized as a 24 kDa proprotein that contains a 36 amino acid (aa) propeptide and a 158 aa mature region (2). Under inflammatory conditions, the propeptide is cleaved by Caspase-1 in the cytoplasm to liberate the mature nonglycosylated 18 kDa monomeric IL-18 (3, 4). Mature rat IL-18 shares 91% aa sequence identity with mouse IL-18 and 60% - 64% aa sequence identity with human, canine, feline, porcine, and rhesus IL-18. IL-18 is secreted by a variety of cell types including macrophages, dendritic cells, and epithelial cells (1, 5). Circulating mature IL-18 is sequestered by soluble IL-18 binding proteins (IL-18 BP) that inhibit IL-18 bioactivity (6). IL-18 interacts with the widely expressed IL-18 R $\alpha$  which then recruits the signaling subunit IL-18 R $\beta$  (7, 8). The IL-1 family member IL-1F7 also binds to IL-18 R $\alpha$  but does not recruit IL-18 R $\beta$  or induce signaling (9). IL-1F7 binds IL-18 BP and enhances its neutralizing effect on IL-18 activity (9). IL-18 synergizes with other cytokines to activate NK, Th1, and Th17 cells and to increase the production of IFN- $\gamma$  (1, 5, 10-12). IL-18 can also promote Th2 cytokine release which reduces the effectiveness of antiviral responses (13, 14). Increased levels of active IL-18 contribute to the severity of autoimmunity and hypertension, while deficiency of IL-18 results in symptoms of metabolic syndrome (1, 5, 15, 16). In cancer, IL-18 stimulates Th1 and NK cells to target tumor cells, but it can also promote angiogenesis, metastasis, and tumor cell immune evasion (11).

**PRODUCT SPECIFIC NOTICES**

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.