

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Semaphorin 4D in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant mouse Semaphorin 3F is observed.
Source	Monoclonal Rat IgG _{2A} Clone # 547406
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Semaphorin 4D Phe24-Thr657 Accession # NP_038688
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Semaphorin 4D (Sema4D/CD100, previously sem J, G or C-like 2) is a widely expressed 150 kDa type I transmembrane glycoprotein of the Class 4 Semaphorin family with activity in the immune and nervous systems. The 861 amino acid (aa) Semaphorin 4D contains a 710 aa extracellular domain (ECD) with a sema domain and an immunoglobulin-like domain. Within the ECD region used as an immunogen, mouse Semaphorin 4D shares 87% and 93% aa identity with human and rat Semaphorin 4D, respectively. A soluble, active 120 kDa form is produced proteolytically upon activation of B and T cells. Semaphorin 4D binds Plexin B1, but also CD72 on B and dendritic cells. In developing neurons, Semaphorin 4D assists in guidance. Semaphorin 4D is proangiogenic when expressed by tumor-associated macrophages.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.