

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Pro EGF in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant mouse (rm) Amphiregulin, rmEpigen, rmEpiregulin, rmEGF, rmMFG-E8, recombinant human (rh) EGF, rhHB-EGF, rhHRG alpha, rhHRG beta, or recom
Source	Monoclonal Rat IgG _{2A} Clone # 463004
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse EGF Trp29-Arg1029 Accession # P01132
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

EGF is the prototypic member of a family of growth factors that also includes amphiregulin, betacellulin, epigen, epiregulin, HB-EGF, neuregulins-1 through -6, and TGF-α (1). These proteins contain EGF-like domains with three intramolecular disulfide bonds between conserved cysteines (2). EGF family members are synthesized as transmembrane preproteins with varying numbers of EGF-like domains (3). The extracellular region of mouse pro-EGF contains eight LDL R class B repeats and nine EGF-like domains (4). Mature EGF is derived from the juxtamembrane EGF-like domain. EGF binds ErbB1 and induces the formation of homodimers or heterodimers containing ErbB2 (5). Pro-EGF is most highly expressed in the submaxillary gland and kidney (6). In the kidney, the 190 kDa preproprotein is cleaved by membrane-associated serine proteases, liberating the extracellular region which is subsequently processed into smaller fragments including the 6 kDa mature EGF (7-10). The various cleavage products produced in the kidney also are present in urine (9-11). In the submaxillary gland, however, nearly all EGF is processed intracellularly and stored in secretory vesicles (6, 12). The soluble precursor binds ErbB1 and induces cellular proliferation, although it is significantly less potent than mature EGF (8, 9). In human thyroid carcinoma cells, a splice variant of pro-EGF with a deletion in the cytoplasmic domain induces increased proliferative activity relative to wild-type pro-EGF (13). Within the extracellular region, mouse pro-EGF shares 79% amino acid sequence identity with rat pro-EGF and 67%-69% with human, canine, feline, and porcine pro-EGF.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.