

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human T-bet/TBX21 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 525831
Purification	Protein A or G purified from cell culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human T-bet/TBX21 Glu326-Asn535 Accession # Q9UL17
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Intracellular Staining by Flow Cytometry	0.25-1 µg/10 ⁶ cells	Human peripheral blood mononuclear cells (PBMCs) treated with Anti-Human IL-4 Polyclonal Antibody (Catalog # AB-204-NA; 20 µg/mL) and Recombinant Human IL-12 (Catalog # 219-IL; 10 ng/mL) for 5 days to induce Th1 cell development were fixed and permeabilized with FlowX FoxP3 Fixation & Permeabilization Buffer Kit (Catalog # FC012)

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

T-box expressed in T cells (T-bet), also known as T-box transcription factor TBX21, is a 62 kDa member of the T-box family of transcription factors and the Tbr1 subfamily. Human T-bet is 535 amino acids in length and contains a T-box DNA binding domain (aa 136-327). Human T-bet shares 88% aa sequence identity with mouse T-bet. T-bet is a nuclear protein highly apparent in Th1 cells. Northern blot analysis revealed that it is also expressed in lung, thymus and spleen. Functionally, T-bet controls the expression of the Th1 cytokine, IFN-γ, and initiates Th1 lineage development from naïve Th precursor cells by both activating Th1 genetic programs and by repressing the opposing Th2 programs.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.