

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse TSLP R in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse TSLP R Ala20-Leu233 Accession # BAA92159
Conjugate	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

TSLP R, also named Delta (1) and CRLM-2 (2) (cytokine receptor-like module-2), was originally cloned as a novel type 1 cytokine receptor with similarity to the common gamma chain. It was subsequently identified to be a subunit of the cellular receptor for the IL-7-like cytokine TSLP and termed TSLP R (3). The TSLP R cDNA encodes a 359 amino acid (aa) residue type 1 membrane protein with a 24 aa residue signal peptide, a 206 aa residue extracellular domain that showed 24% sequence identity with the mouse common γ receptor, a 23 aa residue transmembrane domain, and a 106 aa residue cytoplasmic domain. The cytoplasmic domain of TSLP R contains a membrane-proximal box1 motif which is known to be important for association with JAKs. An alternatively spliced mRNA variant encoding a soluble TSLP R has also been reported (2). TSLP R expression is ubiquitous in the immune and hematopoietic cells, but is up-regulated in Th2-skewed cells. Cells expressing TSLP R alone bind TSLP with low affinity. Co-expression of TSLP R and IL-7 R α is required for high-affinity TSLP binding and signal transduction (3, 4).

References:

1. Fujio, K. *et al.* (2000) *Blood* **95**:2204.
2. Hiroyama, T. *et al.* (2000) *Biochem. Biophys. Res. Commun.* **272**:224.
3. Park, L.S. *et al.* (2000) *J. Exp. Med.* **192**:659.
4. Pandey, A. *et al.* (2000) *Nat. Immunol.* **1**:59.