

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human N-Acetylglucosaminyltransferase V/MGAT5 in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG _{2A} Clone # 706824
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human N-Acetylglucosaminyltransferase V/MGAT5 Leu189-Leu741 Accession # Q09328
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

N-Acetylglucosaminyltransferase V (GnT-V), also known as mannosylglycoprotein N-acetyl-glucosaminyltransferase 5 (MGAT5), adds an N-acetylglucosamine to the α1-6-linked core mannose of an N-linked oligosaccharide in the Golgi apparatus (1). This reaction is the committing step for the biosynthesis of β1-6GlcNAc-branched arm in N-glycans. The degree of N-glycan branching has been shown to regulate cell proliferation and differentiation (2). An increase in the GnT-V activity and its glycan products is also known to positively correlate with the progression of invasive malignancies (3, 4). For example, ectopic expression of GnT-V in epithelial cells results in morphological transformation and tumor growth in mice and overexpression in carcinoma cells has been shown to induce metastatic spread (3-5).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.