

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse IL-15 R α in direct ELISAs.
Source	Monoclonal Rat IgG _{2B} Clone # 888220
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse IL-15 R α Gly33-Lys205 Accession # Q60819
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Interleukin 15 Receptor alpha (IL-15 R α) is a high affinity receptor that specifically binds IL-15 with high affinity and associates as a heterotrimer with the IL-2 receptors beta and gamma subunits to initiate signal transduction. IL-15 R α is expressed on a wide variety of T cells and B cells as well as non-lymphoid cells. IL-15 R α is a 58-60 kDa protein that shares structural similarities to the IL-2 R α protein. IL-15 R α and IL-2 R α genes also share similar intron-exon organization and are closely linked on human chromosome 10p14-p15. Human IL-15 R α shares 45% amino acid (aa) homology with the mouse form of the receptor. Signaling of IL-15 can occur in one of three ways; through the heterotrimeric complex of IL-15 R α , IL-2 R β and IL-2 R γ , through the heterodimeric complex of IL-2 receptors beta and gamma common, and through a novel 60-65 kDa IL-15 RX subunit found on mast cells. The binding of IL-15 to IL-15 R α has been reported to antagonize the TNF- α -mediated apoptosis in fibroblasts by competing with TNFR1 for TRAF2 binding.