

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human GDF-3 in direct ELISAs. In direct ELISAs, less than 10% cross-reactivity with recombinant human (rh) BMP-2 and rhGDF-1 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 878751
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived human recombinant GDF-3/BMP-2 heterodimer Ala251-Gly364 Accession # Q9NR23
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

GDF-3 (previously called Vgr-2) is a TGF- β superfamily member belonging to the growth/differentiation factor family (1, 2). GDF-3 is expressed in undifferentiated embryonic stem (ES) cells, white adipose tissue and the brain (2-4). The 364 amino acid (aa) human GDF-3 contains a 21 aa signal sequence, a 229 aa propeptide and a 114 aa mature region that contains one potential N-glycosylation site. The mature region contains a cysteine-knot structure that is conserved throughout family members. However, it lacks the fourth cysteine which is responsible for the formation of an inter-molecular disulfide bond, so GDF-3 may exist as a non-covalent homodimer (2, 5). Mature human GDF-3 shares 83%, 83%, 91%, 92% and 93% aa sequence identity with mouse, rat, bovine, canine and equine GDF-3, respectively. Most of GDF-3 is present as the uncleaved prepro form (6). The uncleaved and the mature forms both appear to have activity, but that activity may differ (5-8). All forms can oppose BMPs. In human ES cells, inhibition of BMP-2 signaling by GDF-3 maintains pluripotency (5, 7). GDF-3 also influences early cell fate decisions; for example, deletion of mouse GDF-3 produces defects in the anterior visceral endoderm of the pre-gastrulation embryo (6-8). GDF-3 cooperates with GDF-1 in embryogenesis, and the mature protein has nodal-like activity (8, 9). Although GDF family members signal through BMP receptors (ALK-1, -2, -3 and -6), which activate Smads 1, 5 and 8, GDF-3 signaling through ALK-4 and ALK-7, which activate Smads 2 and 3, has also been reported (9, 10). In adipocytes, GDF-3 is induced by a high fat diet, promoting adipogenesis and obesity (3, 10, 11).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.