

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Protocadherin-1 in direct ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 648127
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Protocadherin-1 Thr58-Asn162 Accession # Q08174
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	0.25-1 µg/10 ⁶ cells	A431 human epithelial carcinoma cell line

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

Protocadherin-1 (PCDH-1; also PC42) is a 150 - 170 kDa δ1 subgroup member of the nonclustered protocadherin family of molecules. It is expressed on macrophages, respiratory epithelium, endothelial cells and neurons. Protocadherin-1 apparently forms homophilic Ca²⁺-dependent complexes and likely serves as an adhesion molecule. Human full-length Protocadherin-1 precursor is 1237 amino acids (aa) in length. It is a type I transmembrane glycoprotein that contains a 795 aa extracellular domain (ECD) (aa 58-852) plus a 364 aa cytoplasmic region. There are seven cadherin domains in the ECD (aa 58-844), and an RRVTF cytoplasmic motif that binds PP1α phosphatase. There are multiple splice variants. Alternative start sites exist at Met23 and Met235, there is a deletion of aa 209-220, and a 27 aa substitution for aa 1034-1237. Over aa 58-162, human Protocadherin-1 shares 96% aa identity with mouse Protocadherin-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.