

Human CD98 Fluorescein-conjugated Antibody

Monoclonal Mouse IgG_{2A} Clone # 590559

Catalog Number: FAB5920F


100 Tests

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human CD98 in direct ELISAs.		
Source	Monoclonal Mouse IgG _{2A} Clone # 590559		
Purification	Protein A or G purified from hybridoma culture supernatant		
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human CD98 Asp105-Gln529 Accession # P08195		
Conjugate	Fluorescein Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm (FITC)		
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.		
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μL/10 ⁶ cells	See Below



PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage

Protect from light. Do not freeze.

12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CD98, also known as 4F2hc, is a ubiquitously expressed, 85-95 kDa member of the SLC3 (solute carrier 3) family of amino acid transporters. It is a type II transmembrane glycoprotein that covalently-associates with a variable number of small, 38-45 kDa, 12-transmembrane proteins that belong to the SLC7 family of molecules. The 120-130 kDa heterdimeric 4F2 complex is known to transport neutral or cationic amino acids with, or without, a contribution of sodium. It also interacts with Integrin β1 and β3 to promote cell polarization and migration. Notably, CD98 is "vertebrate-restricted," and its appearance evolutionarily has been linked to the onset of tumorigenesis. Over amino acids (aa) 105-529, human and mouse CD98 share 76% aa sequence identity.

Rev. 2/6/2018 Page 1 of 1

