

## DESCRIPTION


<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human LRP-4 in ELISAs.
<b>Source</b>	Monoclonal Mouse IgG <sub>2A</sub> Clone # 814334
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Chinese hamster ovary cell line CHO-derived recombinant human LRP-4 Ser21-Ser1725 (predicted) Accession # O75096
<b>Conjugate</b>	Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm
<b>Formulation</b>	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	10 $\mu$ L/10 <sup>6</sup> cells	See Below

## DATA



## PREPARATION AND STORAGE

**Shipping** The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

**Stability & Storage** **Protect from light. Do not freeze.**

- 12 months from date of receipt, 2 to 8 °C as supplied.

## BACKGROUND

LRP-4 (Low density lipoprotein-related protein #4; also MEGF7 and LRP13) is a 220-270 kDa glycoprotein, member of the LDLR family of proteins. It is expressed on neurons, oocytes, spermatogonia and skeletal muscle cells, and binds multiple ligands, including WISE, apoE, MuSK and neuronal Agrin. It serves to negatively regulate Wnt signaling during development, and to cluster AChRs at neuromuscular junctions. Mature human LRP-4 is an 1885 amino acid (aa) type I transmembrane glycoprotein. It contains a 1705 aa extracellular domain (ECD) (aa 21-1725) plus a 159 aa cytoplasmic region (aa 1747-1905). In the ECD, there are eight LDLR class A repeats (aa 26-350), two EGF-like repeats (aa 354-434) and 20 LDLR class B repeats that contain an intervening EGF-like domain (aa 480-1610). Over aa 18-344, human LRP-4 shares 88% aa sequence identity with mouse LRP-4.