

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse IL-21 R in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human IL-21 R or recombinant mouse IL-2 Rβ is observed.
Source	Monoclonal Rat IgG ₁ Clone # 155516
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant mouse IL-21 R Cys20-Pro236 Accession # Q9JHX3
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-21 R (interleukin-21 receptor) is a type I transmembrane glycoprotein within the class I cytokine receptor family, type 4 subfamily (1-5). Complex formation between IL-21 R and the common γ chain (γ_c), also used for IL-2, IL-4, IL-7, IL-9, IL-13 and IL-15 receptors, is required for signaling (6, 7). Mouse IL-21 R cDNA encodes 521 amino acid (aa) including a 19 aa signal peptide, a 218 aa extracellular domain (ECD) with 4 conserved cysteine residues, a fibronectin type III domain, and a WSXWS motif, a 21 aa transmembrane domain and a 271 aa cytoplasmic domain with a Box 1 motif, a kinase domain, and several sites for tyrosine phosphorylation (4, 5). One such site, pY510, mediates STAT binding (1, 2). The mouse IL-21 R ECD shares 69%, 91%, 65%, 63% and 58% aa identity with human, rat, equine, canine and bovine IL-21 R, respectively. One potential 447 aa isoform, with an alternate start site at aa 83, lacks the four conserved ECD cysteines. IL-21 R is expressed mainly on B cells (highest on mature, activated, follicular and germinal center B cells), NK cells, and activated T cells, but is also found on dendritic cells, alternatively activated macrophages, intestinal lamina propria fibroblasts and epithelial cells, and keratinocytes (1, 3-5). Both IL-21 and IL-4 are necessary for efficient B cell IgG1 production and normal germinal center architecture (8). B cell IL-21 R engagement induces Blimp-1 (which mediates plasma cell differentiation), and is important for memory responses (1, 9, 10). IL-21 R engagement on mouse NK cells enhances their cytotoxic activity and IFN-γ production (4, 11). IL-21 R engagement on CD8⁺ T cells aids control of viral infection and tumor growth; IL-21 R is also necessary for sufficient numbers of regulatory T cells to combat chronic inflammation (1, 12, 13). IL-21 R expression is often upregulated in allergic skin inflammation, systemic lupus erythematosus and diffuse large B cell lymphoma (DLBCL) (1, 2, 14, 15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.