

## DESCRIPTION

<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse CCR1 in direct ELISAs.
<b>Source</b>	Recombinant Monoclonal Rat IgG <sub>2B</sub> Clone # 643854R
<b>Purification</b>	Protein A or G purified from cell culture supernatant
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant mouse CCR1 Extracellular regions (aa 1-34, aa 92-107, aa 172-192, aa 265-281) Accession # P51675
<b>Conjugate</b>	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Flow Cytometry</b>	0.25-1 µg/10 <sup>6</sup> cells	RAW 264.7 mouse monocyte/macrophage cell line

## PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

## BACKGROUND

CCR1 (C-C chemokine receptor 1; also MIP-1a receptor and CD191) is a 43 kDa member of the GPCR #1 family of transmembrane proteins. Mouse CCR1 is expressed on osteoclasts, IL-13<sup>+</sup> T cells, neutrophils, bone marrow-derived mast cells, eosinophils, monocytes and vascular intimal smooth muscle cells. Multiple chemokines are reported to bind to CCR1. MIP-1a/CCL3 and RANTES/CCL5 are natural ligands, while CCL15, CCL9 and CCL23 are potent antagonists after N-terminal processing. Mouse CCR1 is a 7-transmembrane protein that is 355 amino acids (aa) in length. It contains a 34 aa N-terminal extracellular domain plus a 50 aa C-terminal cytoplasmic tail. Over aa sequences 1-34, 92-107, 172-192 and 265-281 collectively, mouse CCR1 shares 76% and 89% aa identity with human and rat CCR1, respectively.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.