

DESCRIPTION

| | |
|---------------------------|--|
| Species Reactivity | Mouse |
| Specificity | Detects mouse LIF R α in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human LIF R α is observed. |
| Source | Monoclonal Rat IgG ₁ Clone # 673602 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant mouse LIF R α Leu44-Ser828 Accession # P42703 |
| Conjugate | Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm |
| Formulation | Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

| | Recommended Concentration | Sample |
|-----------------------|--------------------------------------|-----------------------------------|
| Flow Cytometry | 0.25-1 μ g/10 ⁶ cells | D3 mouse embryonic stem cell line |

PREPARATION AND STORAGE

| | |
|--------------------------------|--|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

Leukemia Inhibitory Factor Receptor alpha (LIF R α), also known as LIFR beta and CD118, is a 190 kDa type I transmembrane protein in the Interleukin-6 receptor family. Members of this family mediate the biological effects of Cardiotrophin-1, CLC, CNTF, IL-6, IL-11, IL-27, and Oncostatin M (1). Mature mouse LIF R α consists of a 785 amino acid (aa) extracellular domain (ECD) with two cytokine receptor homology domains, one WSxWS motif, and three fibronectin type III repeats, followed by a 25 aa transmembrane segment and a 239 aa cytoplasmic domain (2, 3). Within the ECD, mouse LIF R α shares 73% and 90% aa sequence identity with human and rat LIF R α , respectively. Alternative splicing generates a 90 kDa soluble form of the mouse LIF R α ECD (4). LIF R α binds the pleiotropic cytokine LIF with low affinity, and the soluble isoform retains LIF-binding activity (5). Binding affinity is increased by the ligand-induced association of LIF R α with the signal transducing subunit gp130 (6, 7). The LIF R α /gp130 receptor complex also transduces Oncostatin M signals, although LIF R α alone does not interact with Oncostatin M (6). gp130 associates with different ligand-specific receptors to form signaling receptor complexes for the other IL-6 family ligands (1). The CNTF receptor is a ternary complex that contains CNTF R α and gp130 as well as LIF R α (8, 9). LIF R α is widely expressed, and LIF induces the proliferation, differentiation, and activation of cells in many tissues (10, 11). In particular, LIF R α plays an important role in several aspects of early pregnancy such as blastocyst implantation in the uterus (4, 12-14).

References:

- Muller-Newen, G. (2003) Science STKE **2003**:pe40.
- Gearing, D.P. *et al.* (1991) EMBO J. **10**:2839.
- Tomida, M. *et al.* (1994) J. Biochem. **115**:557.
- Owczarek, C.M. *et al.* (1996) J. Biol. Chem. **271**:5495.
- Layton, M.J. *et al.* (1992) Proc. Natl. Acad. Sci. **89**:8616.
- Gearing, D.P. *et al.* (1992) Science **255**:1434.
- Giese, B. *et al.* (2005) J. Cell Sci. **118**:5129.
- Ip, N.Y. *et al.* (1992) Cell **69**:1121.
- Davis, S. *et al.* (1993) Science **260**:1805.
- Metcalf, D (2003) Stem Cells **21**:5.
- Kubota, Y. *et al.* (2008) J. Clin. Invest. **118**:2393.
- Paiva, P. *et al.* (2009) Cytokine Growth Factor Rev. **20**:319.
- Stewart, C.L. *et al.* (1992) Nature **359**:76.
- Cheng, J.-G. *et al.* (2001) Proc. Natl. Acad. Sci. **98**:8680.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.