

#### DESCRIPTION

<b>Species Reactivity</b>	Human
<b>Specificity</b>	Detects human CLEC9a in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) CLEC1, 2, 2A, 3B, 10A, 12B, 14A, rhCD302/CLEC13a, rhM1CL, or recombinant mouse CLEC9a is observed.
<b>Source</b>	Monoclonal Mouse IgG <sub>1</sub> Clone # 683409
<b>Purification</b>	Protein A or G purified from hybridoma culture supernatant
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant human CLEC9a Lys57-Val241 Accession # Q6UXN8
<b>Conjugate</b>	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
<b>Formulation</b>	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details.  *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

#### APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
<b>Flow Cytometry</b>	0.25-1 µg/10 <sup>6</sup> cells	Human HEK293 cells transfected with human CLEC9a

#### PREPARATION AND STORAGE

<b>Shipping</b>	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
<b>Stability &amp; Storage</b>	<b>Protect from light. Do not freeze.</b> <ul style="list-style-type: none"> <li>12 months from date of receipt, 2 to 8 °C as supplied.</li> </ul>

#### BACKGROUND

CLEC9a (C-type lectin domain family 9 member A), also known as DNGR-1, is a type II transmembrane glycoprotein member of the C-type lectin superfamily (1, 2). Mature human CLEC9a consists of a 35 amino acid (aa) cytoplasmic domain with an ITAM-like motif, a 21 aa transmembrane segment, and a 185 extracellular domain (ECD) that contains a stalk region and one C-type lectin domain (CTLD) (3-5). Within the ECD, human CLEC9a shares 57% aa sequence identity with mouse and rat CLEC9a. Although the CTLD of CLEC9a structurally resembles that of other C-type lectins, it lacks the conserved residues that typically mediate calcium and carbohydrate binding. CLEC9a is expressed as a disulfide-linked homodimer of approximately 50 kDa N-glycosylated subunits (3, 5). Human CLEC9a expression is restricted to a subpopulation of BDCA-3<sup>+</sup> conventional dendritic cells (cDC) and CD16<sup>+</sup> monocytes (3-5). BDCA-3<sup>+</sup> cDC are analogous to mouse CD8<sup>+</sup> cDC which are specialized in antigenic cross-presentation in complex with MHC class I molecules (6). In mouse, CLEC9a is additionally expressed on plasmacytoid dendritic cells (4, 5). CLEC9a ligation enhances antigen uptake and processing, leading to presentation on MHC class I and cytotoxic T cell (CTL) priming (3-5). In mouse, CLEC9a recognizes normally intracellular determinant(s) of necrotic cells and mediates their uptake by the dendritic cell (7). The subsequent antigenic cross-presentation to CTL is important for clearing necrotic cellular debris (7). CLEC9a signaling triggers activation of the tyrosine kinase Syk (3, 7).

#### References:

- Huysamen, C. and G.D. Brown (2009) *FEMS Microbiol. Lett.* **290**:121.
- Geijtenbeek, T.B.H. *et al.* (2004) *Annu. Rev. Immunol.* **22**:33.
- Huysamen, C. *et al.* (2008) *J. Biol. Chem.* **283**:16693.
- Caminschi, I. *et al.* (2008) *Blood* **112**:3264.
- Sancho, D. *et al.* (2008) *J. Clin. Invest.* **118**:2098.
- Dudziak, D. *et al.* (2007) *Science* **315**:107.
- Sancho, D. *et al.* (2009) *Nature* **458**:899.

#### PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.