

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TCF7L1 in direct ELISAs. In direct ELISAs, 5-10% cross-reactivity with recombinant human (rh) TCF-4 (aa 102-516) and rhTCF-12 is observed. No cross-reactivity with rhTCF7L1 (aa 151-320), rhTCF7L2, rhTCF-1, -2, -3, -4 (aa 241-365), rhLEF-1, o
Source	Monoclonal Mouse IgG _{2B} Clone # 622782
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E.coli</i> -derived recombinant human TCF7L1 Lys429-Ser581 Accession # Q9HCS4
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Immunohistochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Transcription factor 7-like 1 (TCF7L1; also HMG box transcription factor 3, TCF-3) is a 63 kDa member of the TCF/LEF family of proteins. Human TCF7L1 is 588 amino acids (aa) in length and contains one HMG box DNA-binding domain (aa 346-414). Human TCF7L1 shares 95% aa sequence identity with mouse TCF7L1. TCF7L1 is expressed in hair follicles and skin keratinocytes and at lower levels in stomach epithelium. Functionally, it participates in the Wnt signaling pathway. TCF7L1 binds to DNA and acts as a repressor in the absence of CTNNB1 and as an activator in its presence. TCF7L1 is also necessary for the terminal differentiation of epidermal cells, the formation of keratohyalin granules, and the development of the barrier function of the epidermis.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.