

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Semaphorin 4C in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human Semaphorin 3A, 3B, 3C, 3D, 3E, 3F, 4A, 4B, 4D, 4G, 5A, 5B, 6A, 6B, 6C, 6D, 7A, recombinant mouse Semaphorin 3B, 3C, 3E, 3F, 4A, 4B, 4C, 4D, 4F, 4G,
Source	Monoclonal Mouse IgG ₁ Clone # 660016
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Semaphorin 4C Ala21-Gly663 (predicted) Accession # Q9C0C4
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Semaphorin 4C (Sema4C; also Sema I and M-SemaF) is a 100-105 kDa, class IV member of the semaphorin family of proteins. It is expressed by precursors to neurons and myocytes, and may regulate their differentiation into mature forms. Mature human Semaphorin 4C is a type I transmembrane glycoprotein that is 813 amino acids (aa) in length. It contains a 643 aa extracellular region (aa 21-663) that is characterized by the presence of one Sema domain (aa 30-497), a PSI region (aa 499-551), and an Ig-like C2-type domain (aa 556-644). The cytoplasmic region interacts with PZD-domain containing proteins. There are three potential splice variants. One demonstrates an alternative start site at Met324, a second shows a deletion of aa 173-211, and a third contains an 89 aa substitution for aa 1-36. Over aa 21-663, human Semaphorin 4C shares 85% aa identity with mouse Semaphorin 4C.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.