

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Proteinase 3/Myeloblastin/PRTN3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human Granzyme B is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 684042
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf21-derived recombinant human Proteinase 3/Myeloblastin/PRTN3 Ala26-Arg249 Accession # P24158
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Proteinase 3 (PRTN3), also known as Myeloblastin (MBN), is a 32-33 kDa member of the peptidase S1 family of enzymes. It is expressed by monocytes and neutrophils, the latter of which either secretes it, sequesters it in azurophilic granules, or expresses it on the cell surface. When secreted, it acts on HK and activates the kinin pathway. In azurophilic granules, it aids in the digestion of phagocytosed material. On the cell surface, it likely acts on ECM. Human PRTN3 precursor is 231 amino acids (aa) in length. It contains an Ala26Glu27 propeptide that is removed during maturation, a 221 aa mature enzyme (aa 28-248), and an eight aa C-terminal propeptide (aa 249-256). Within the cell, a 35 kDa immature form exists; on the cell surface, both constitutively inactive, and induced active forms may be found, often in a noncovalent association with CD177/NB1. Over aa 26-249, human PRTN3 shares 68% aa identity with mouse PRTN3.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.