

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human TRIM21 in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) TRIM5 or rhTRIM32 is observed.
Source	Monoclonal Mouse IgG _{2A} Clone # 672714
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human TRIM21 Arg195-Pro293 Accession # P19474
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TRIM21 (Tripartite motif-containing protein 21; also Ro(SS-A), 52 kDa Ro Protein/Ro52, and RING finger protein 81) is a 52-56 kDa member of the RING finger-B-box-coiled-coil family of proteins. It is an E3 ligase that is found in both nucleus and cytoplasm, where it is often associated with microtubules. TRIM21 ubiquitinates select proteins. In B cells, it targets the Fc fragment of misfolded IgG, providing QC on its production. In macrophages, it acts in a nondegradative manner on IRF8, promoting innate immunity. Human TRIM21 is 475 amino acids (aa) in length. It contains one E3 ligase RING finger domain (aa 16-55), a B-Box type zinc-finger region (aa 92-123), a coiled-coil region (aa 128-238) and a C-terminal SPRY/B30.2 Ig binding domain (aa 268-465). TRIM21 is reported to form trimers. Over aa 195-293, human TRIM21 exhibits 72% aa identity with mouse TRIM21.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.