

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human TRANCE/TNFSF11/RANK L in direct ELISAs and Western blots. In direct ELISAs and Western blots, 100% cross-reactivity with recombinant mouse (rm) TRANCE and approximately 25% cross-reactivity with recombinant human (rh) LIGHT is observed. No c
Source	Monoclonal Mouse IgG ₃ Clone # 685809
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human TRANCE/TNFSF11/RANK L Gly64-Asp245 Accession # AAC51762
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

TNF-related activation-induced cytokine (TRANCE; also RANKL, OPGL, and ODF) is a 35 kDa (predicted) type II transmembrane glycoprotein and member of the TNF cytokine family. Human TRANCE is 317 amino acids (aa) in length and contains a 47 aa cytoplasmic region, a 21 aa transmembrane region, and a 249 extracellular domain (ECD), which contains two potential sites of N-linked glycosylation. Splicing variants produce three isoforms for human TRANCE. Isoform 1 is the standard form. In isoform 2, aa corresponding to 1-73 in isoform 1 are missing. In isoform 3, aa 1-47 are missing. Human TRANCE ECD is 83% identical to mouse TRANCE ECD. TRANCE is expressed highest in the peripheral lymph nodes and weaker in the spleen, peripheral blood leukocytes, bone marrow, heart, placenta, skeletal muscle, stomach, and thyroid. TRANCE plays a role in osteoclast differentiation and activation.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.