

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human JADE-1/PHF-17 in direct ELISAs and Western blots. In Western blots, approximately 50% cross-reactivity with recombinant human Jade-2/PHF15 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 658005
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human JADE-1/PHF-17 Met1-Ser137 Accession # Q6IE81
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Jade-1 (Gene for Apoptosis and Differentiation in Epithelia; also PHF17) is a 94 kDa member of the JADE family of proteins. It is expressed in renal tubule cells as well as embryonic primitive streak (in mouse). Initially, Jade-1 was described as a proapoptotic tumor suppressor. Jade-1 also interacts with the HBO1 complex to specifically acetylate nucleosome histones, thus acting as a transcriptional activator that likely impacts DNA replication, recombination and repair. Human Jade-1 is 842 amino acids (aa) in length. It contains one PEST domain with multiple Ser phosphorylation sites (aa 5-28), and two PHD Zinc fingers (aa 203-253 and 312-371) that are required for histone acetylation. There are two splice variants, one that is 61-64 kDa in size and contains a 7 aa substitution for aa 503-842, and another that shows a deletion of aa 99-110. Over aa 1-137, human Jade-1 shares 96% aa identity with mouse Jade-1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.