

Human Lymphotoxin β R/TNFRSF3 PE-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 71319

Catalog Number: FAB629P
100 Tests

DESCRIPTION


| | |
|---------------------------|---|
| Species Reactivity | Human |
| Specificity | Detects human Lymphotoxin β R/TNFRSF3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, no cross-reactivity with recombinant human (rh) 4-1BB, rhBAFF R, rhCD27, rhCD30, rhCD40, rhDR3, rhDR6, rhEDAR, rhFas, rhGITR, rhHVEM, recombinant mouse (rm) Lymphotoxin β R, rhNGF R, rhOPG, rhRELT, rhTAJ, rhTNF RI or rhTNF RII is observed. |
| Source | Monoclonal Mouse IgG ₁ Clone # 71319 |
| Purification | Protein A or G purified from hybridoma culture supernatant |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant human Lymphotoxin β R/TNFRSF3 Gln31-Met227 Accession # P36941 |
| Conjugate | Phycoerythrin Excitation Wavelength: 488 nm Emission Wavelength: 565-605 nm |
| Formulation | Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

| | Recommended Concentration | Sample |
|-----------------------|----------------------------------|---------------|
| Flow Cytometry | 10 μ L/10 ⁶ cells | See Below |

DATA



PREPARATION AND STORAGE

| | |
|--------------------------------|---|
| Shipping | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| Stability & Storage | Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied. |

BACKGROUND

Lymphotoxin beta Receptor (LT β R), also known as TNF RIII and TNF R-Related Protein (TNF Rrp) is a member of the TNF receptor superfamily, designated TNFRSF3. Human LT β R cDNA encodes a 435 amino acid (aa) residue type I membrane protein with a putative 30 aa residue signal peptide, a 193 aa residue extracellular domain and a 171 aa residue cytoplasmic domain. The extracellular domain of LT β R contains four cysteine-rich motifs characteristic of the TNF receptor superfamily. The cytoplasmic region of LT β R shares little sequence similarity with other TNF receptor family members, suggesting that different signaling mechanisms may be used. LT β R is expressed in a variety of tissues including visceral and lymphoid tissues. LT β R is also expressed by cell lines of monocytic, epithelial, and fibroblastic origins but not by T and B lymphocytes. Human and mouse LT β R share 76% aa sequence homology. The TNF family ligands that have been shown to bind and activate LT β R include LIGHT (also a ligand for HVEM) and the heterotrimeric Lymphotoxin LT α 1/ β 2 or LT α 2/ β 1. Depending on the cell type, activation of LT β R has been shown to induce NF κ B activation, chemokine production, growth arrest, and apoptosis. *In vivo*, LT β R has been shown to play a critical role in controlling cellular immune functions and lymphoid organogenesis.

References:

1. Zhai, Y. *et al.* (1998) J. Clin. Invest. **102**:1142.
2. Rennert, P.D. *et al.* (1998) Immunity **9**:71.
3. Degli-Esposti, M.A. *et al.* (1997) J. Immunol **158**:1756.
4. Mackay, F. *et al.* (1996) J. Biol. Chem. **271**:8618.
5. Crowe, P.D. *et al.* (1994) Science **264**:707.