

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Heparan Sulfate 2-O-Sulfotransferase 1/HS2ST1 in ELISAs and Western blots.
Source	Monoclonal Mouse IgG ₁ Clone # 738306
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Heparan Sulfate 2-O-Sulfotransferase 1/HS2ST1 Met59-Asn356 Accession # Q7LGA3
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Heparan sulfate is a highly sulfated polysaccharide found on the cell surface and within the extracellular matrix. Typically, it is covalently attached to the protein core of proteoglycans, such as syndecans and glypicans. Heparin, on the other hand, can be considered as a highly sulfated version of heparan sulfate that is predominantly found in mast cells. Both heparin and heparan sulfate contain disaccharide repeats of uronic acid and N-acetylglucosamine and are modified by the same sulfotransferases (1, 2). The uronic acid residues are either glucuronic acid or iduronic acid and maybe sulfated at the 2-O position by heparan sulfate 2-O sulfotransferase 1 (HS2ST1) (3, 4). HS2ST1 physically interacts in the Golgi apparatus with glucuronyl c5-epimerase (5), which catalyzes the conversion of glucuronic acid to iduronic acid (6). As a consequence, 2-O sulfation predominantly occurs on iduronic acids naturally and overexpression of HS2ST1 alone causes an increase in 2-O sulfation on glucuronic acid (7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.