

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human LILRA-2 in direct ELISAs. In direct ELISAs, no cross-reactivity with recombinant human (rh) LILRA-1, rhLILRA-3, rhLILRA-4, rhLILRA-5, rhLILRA-6, rhILT-2, rhILT-4, or rhILT-5 is observed.
Source	Monoclonal Mouse IgG _{2B} Clone # 600002
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LILRA2/CD85h/ILT1 Pro17-Ser437 Accession # Q8N149
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Immune-like transcript 1 (ILT1), also known as leukocyte immunoglobulin-like receptor 7 (LIR-7), leukocyte immunoglobulin-like receptor subfamily A member 2 (LILRA2), and CD85H, is a 51 kDa (unglycosylated) type I transmembrane glycoprotein and member of the leukocyte immunoglobulin-like receptor family. Human ILT1 is synthesized as a 483 amino acid (aa) precursor that contains a 23 aa signal sequence, a 426 aa extracellular domain (ECD), a 21 aa transmembrane segment, and a 13 aa cytoplasmic region. The ECD contains four Ig-like C2-type domains and seven potential sites for N-linked glycosylation. A splice variant produces a second isoform that has a one aa substitution for aa 419-436 in the longer form. There are no murine orthologs for human ILT1. ILT1 may act as a receptor for class I MHC antigens.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.