

TGF-β3 Alexa Fluor® 488-conjugated Antibody

Monoclonal Mouse IgG₁ Clone # 44922 Catalog Number: FAB643G

100 µg

DESCRIPTION		
Specificity	Detects TGF-β3 from multiple species in ELISAs and Western blots. In ELISAs, does not cross-react with recombinant human (rh) TGF-β1or rhTGF-β2.	
Source	Monoclonal Mouse IgG ₁ Clone # 44922	
Purification	Protein A or G purified from hybridoma culture supernatant	
Immunogen	S. <i>frugiperda</i> insect ovarian cell line <i>Sf</i> 21-derived recombinant human TGF-β3 Ala301-Ser412 (Tyr340Phe) Accession # NP_003230.1	
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm	
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS				
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.				
ELISA Capture (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.			
ELISA Detection (Matched Antibody Pair)	Optimal dilution of this antibody should be experimentally determined.			
Western Blot	Optimal dilution of this antibody should be experimentally determined.			
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.			

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied	

BACKGROUND

The active form of TGF- β 3 is a disulfide-linked homodimer which is generated by activation of the latent TGF- β 3 complex. TGF- β 5 are multifunctional cytokines that activate or repress multiple genes in a wide variety of cell types. Depending on the cell context, TGF- β 6 can control many cellular processes including growth, morphogenesis, and apoptosis.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/22/2025 Page 1 of 1

China | info.cn@bio-techne.com TEL: 400.821.3475