

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse CEACAM-1/CD66a in direct ELISAs and Western blots. In direct ELISAs, no cross-reactivity with recombinant human (rh) CEACAM-1, -3, -5, or -6 is observed. In Western blots, no cross-reactivity with rhCEACAM-1, -3, -4, -5, -6, or -7 is observed.
Source	Monoclonal Rat IgG ₁ Clone # 723629
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse CEACAM-1/CD66a Glu35-Gly428 Accession # P31809
Conjugate	Allophycocyanin Excitation Wavelength: 620-650 nm Emission Wavelength: 660-670 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	10 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

CEACAM-1 (Carcinoembryonic Antigen-related Cell Adhesion Molecule 1), also known as BGP-1, CD66a and MHVR1, is a 110-120 kDa member of the CEACAM subfamily, CEA family of proteins. It has a wide expression pattern, being found on neutrophils, dendritic cells, endothelial cells, colonic epithelium and hepatocytes. It mediates cell adhesion, and appears to regulate insulin levels and signaling by interacting with the insulin receptor. It also demonstrates proangiogenic effects by inducing endothelial cells to proliferate and form capillary-like tubules. Finally, CEACAM-1 is a known receptor for mouse hepatitis virus. Mature mouse CEACAM-1 is a 487 amino acid (aa) type I transmembrane glycoprotein. Its contains a 394 aa extracellular region (aa 35-428) that shows one V-type (aa 35-142) and three C2-type (aa 147-411) Ig-like domains, plus a 74 aa cytoplasmic domain. Three alternate splice forms exist. One contains a four aa substitution for aa 455-521, a second shows a Gln substitution for aa 142-322, and a third possesses a combination of the first two splice patterns. CEACAM-1 forms homodimers. Over aa 35-428, mouse CEACAM-1 shares 56% and 70% aa identity with human and rat CEACAM-1, respectively.