

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human PILR- α in direct ELISAs.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human PILR- α Gln20-Thr196 Accession # Q9UKJ1
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied in a saline solution containing BSA and Sodium Azide. See Certificate of Analysis for details. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Flow Cytometry	5 μ L/10 ⁶ cells	See Below

DATA



PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. <ul style="list-style-type: none"> ● 12 months from date of receipt, 2 to 8 °C as supplied.

BACKGROUND

PILR- α (Paired Immunoglobulin-like, type 2 Receptor alpha) is a monomeric 44-50 kDa type I transmembrane (TM) paired receptor glycoprotein that belongs to the Ig Superfamily. It is expressed by neutrophils, monocytes, macrophages, CD14⁺CD11a⁻ DC, corneal epithelium and retinal pigment cells. It is known to bind to CD99, PANP, Collectin-12, and NPDC-1. The target for PILR- α appears to be sialylated O-glycans. PILR- α also acts as a receptor for HSV and serves as a negative immunomodulator that contains an ITIM. Mature human PILR- α is 284 amino acids (aa) in length. It contains one V-type Ig-like domain in its extracellular region (aa 32-150), and two ITIMs in its cytoplasmic domain (aa 267-272 and 296-301). There are multiple potential splice variants. One is TM and possesses a 35 aa substitution for aa 264-303, while others are soluble, and may either show a deletion of aa 152-224 that may be coupled to the 35 aa substitution noted above, or simply exhibit a 24 aa substitution for aa 152-303. Over aa 20-196, human PILR- α shares only 42% aa identity with mouse PILR- α , and 89% aa identity with human PILR- β .

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.