

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Glucuronosyltransferase 1A1/UGT1A1 in ELISAs.
Source	Monoclonal Mouse IgG ₁ Clone # 856754
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human Glucuronosyltransferase 1A1/UGT1A1 Leu60-Thr168 Accession # P22309
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

UGT1A1 (UDP-glucuronosyltransferase 1-A1; also UDPGT 1-1 and HUG-BR1) is a 52-57 kDa member of the UGT1A subfamily, UGT family of enzymes. UGT1A1 is expressed by the liver, and catalyzes the conjugation of glucuronic acid (GA) from UDPGA to lipophilic acceptors such as (anti-cancer) drugs and bilirubin. Addition of glucuronic acid increases target solubility and facilitates elimination. Mature human UGT1A1 is a 508 amino acid (aa) type I transmembrane ER glycoprotein. It contains a 465 aa luminal domain (aa 26-490) plus a 26 aa cytoplasmic region. The luminal domain is unusual in that aa 157-176 are embedded in the ER membrane. Amino acids 29-444 contain the enzyme active site. The cytoplasmic tail appears to mediate noncovalent homodimerization, and heterodimerization with UGT2B. The signal sequence (aa 1-25) is normally cleaved, but a Lys15Arg mutation blocks insertion into the ER membrane. There are multiple point mutations that impact enzyme activity. One potential splice form is reported that shows a six aa substitution for aa 289-533. Over aa 60-186, human UGT1A1 shares 64% aa identity with mouse UGT1A1.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.