

Human NFATC2 Alexa Fluor® 700-conjugated Antibody

Monoclonal Mouse IgG_{2B} Clone # 639402

Catalog Number: FAB6499N

100 µg

DESCRIPTION					
Species Reactivity	Human				
Specificity	Detects human NFATC2 in direct ELISAs and Western blots. In Western blots, no cross-reactivity with recombinant human NFATC1 or C3 is observed.				
Source	Monoclonal Mouse IgG _{2B} Clone # 639402				
Purification	Protein A or G purified from hybridoma culture supernatant				
Immunogen	E. coli-derived recombinant human NFATC-2 His575-Pro679 Accession # Q13469				
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm				
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide				
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.				

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

China | info.cn@bio-techne.com TEL: 400.821.3475

				AGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.			
Stability & Storage	ability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied			

BACKGROUND

NFATC2 (Nuclear factor of activated T cells C2; also NFAT1 or NFAT2P) is a 135 kDa member of the NFAT family of transcription factors. NFATC2 is found in T cells and mast cells where it regulates cytokine transcription, Th2 cell differentiation, and cell cycle entry. The transactivation function of NFATC2 is regulated by phosphorylation at Ser53, Ser56, Thr116, and Ser170. Human NFATC2 is 925 amino acids (aa) in length. It contains a calcineurin-binding site (aa 111-116), a transactivation domain (aa 119-199), two NLS (aa 251-253 and 664-666), an RHD that binds DNA (aa 392-574), and one NES (aa 904-913). An alternate splice form has a substitution of the C-terminal 18 amino acids. Human NFATC2 shares 96% aa identity with mouse and rat NFATC2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/22/2025 Page 1 of 1