

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human RLK/TKX in direct ELISAs and Western blots.
Source	Monoclonal Mouse IgG _{2B} Clone # 674801
Purification	Protein A or G purified from hybridoma culture supernatant
Immunogen	<i>E. coli</i> -derived recombinant human RLK/TKX Trp150-Val246 Accession # P42681
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

RLK (resting lymphocyte kinase), also called TKX (T cell-expressed kinase) or PTK4, is a protein tyrosine kinase of the TEK family. It is expressed in T cells and some myeloid cells, especially in Th1 cells in inflammatory diseases such as rheumatoid arthritis and Behcet's disease. Upon TCR stimulation, it translocates to the nucleus and induces IFN-γ production. An alternate start site creates a resident nuclear 55 kDa form in addition to the larger (~58 kDa) form that is cytoplasmic or targeted to the plasma membrane via palmitoylation of a cysteine string motif. The region used as an immunogen is common to both forms and shares 79% and 81% amino acid identity with mouse and rat RLK, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.